

Lecture 2: Stochastic Process, Brownian Motion.

Zhongjian Wang*

Abstract

Summary of Stochastic process and Brownian motion

1 Stochastic Processes

Sequence of r.v.'s $X_1, X_2, \dots, X_n, \dots$ occurring at discrete times $t_1 < t_2 < \dots < t_n < \dots$ is called a discrete stochastic process, with joint distribution $F_{X_{i_1}, X_{i_2}, \dots, X_{i_j}}, i_j = 1, 2, \dots$ as its probability law.

Gaussian Process: all joint distributions are Gaussian.

Continuous Stochastic Process: $X(t) = X(t, \omega), t \in [0, 1]$ or $[0, \infty)$, over probability space (Ω, A, P) , is a function of two variables, $X : [0, 1] \times \Omega \rightarrow R$, where X is a r.v. for each t , for each ω , we have a sample path (a realization) or trajectory of the process.

- Quantities on time variability: $\mu(t) = E(X(t)), \sigma^2(t) = Var(X(t))$, covariance:

$$C(s, t) = E((X(s) - \mu(s))(X(t) - \mu(t))),$$

for $s \neq t$.

- Process with independent increment: $X(t_{j+1}) - X(t_j), j = 0, 1, 2, \dots$.

Standard Wiener Process (Brownian Motion): Gaussian process $W(t), t \geq 0$, with independent increment, and:

$$W(0) = 0 \text{ w.p.1, } E(W(t)) = 0, \text{ } Var(W(t) - W(s)) = t - s,$$

for all $s \in [0, t]$.

B.M. Covariance: $C(s, t) = \min(s, t)$.

Stationary Process: all joint distributions are translation (along time) invariant.

Ornstein-Uhlenbeck Process: Gaussian process with $X(0)$ unit Gaussian, $E(X(t)) = 0$, covariance $E(X_s X_t) = e^{-\gamma|t-s|}$ for $s, t \in R, \gamma > 0$.

B.M. Covariance: $C(s, t) = \min(s, t)$, not stationary. O-U stationary.

*Department of Statistics, University of Chicago

2 Diffusion Process

Suppose joint distribution of $X(t)$ has density $p(t_1, x_1; t_2, x_2; \dots; t_k, x_k)$, define conditional probability:

$$P(X(t_{n+1}) \in B | X(t_i) = x_i, i = 1 : n) = \frac{\int_B p(t_1, x_1, \dots, t_n, x_n; t_{n+1}, y) dy}{\int p(t_1, x_1, \dots, t_n, x_n; t_{n+1}, y) dy}$$

for B any Borel set of R .

• *Markov Process* if:

$$P(X(t_{n+1}) \in B | X(t_i) = x_i, i = 1 : n) = P(X(t_{n+1}) \in B | X(t_n) = x_n).$$

Transition probability:

$$P(s, x; t, B) = \int_B p(s, x; t, y) dy,$$

p transition density.

Wiener:

$$p(s, x; t, y) = \frac{1}{\sqrt{2\pi(t-s)}} \exp\left\{-\frac{(y-x)^2}{2(t-s)}\right\},$$

O-U: ($\gamma = 1$)

$$p(s, x; t, y) = \frac{1}{\sqrt{2\pi(1-e^{-2(t-s)})}} \exp\left\{-\frac{(y-xe^{-(t-s)})^2}{2(1-e^{-2(t-s)})}\right\},$$

Chapman-Kolmogorov (C-K) equation:

$$p(s, x; t, y) = \int_{R^1} p(s, x; \tau, z) p(\tau, z; t, y) dz,$$

for $s \leq \tau \leq t$.

The transition density of Wiener obeys equations:

$$p_t = \frac{1}{2} p_{yy}, \quad (s, x) \text{ fixed}$$

forward equation,

$$p_s = -\frac{1}{2} p_{xx}, \quad (t, y) \text{ fixed}$$

backward equation.

• Markov process with transition density is called *diffusion process* if the following limits exist:

Jump:

$$\lim_{t \rightarrow s^+} \frac{1}{t-s} \int_{|y-x| > \epsilon} p(s, x; t, y) dy = 0,$$

Drift:

$$\lim_{t \rightarrow s^+} \frac{1}{t-s} \int_{|y-x| \leq \epsilon} (y-x) p(s, x; t, y) dy = a(s, x),$$

Diffusion:

$$\lim_{t \rightarrow s^+} \frac{1}{t-s} \int_{|y-x| \leq \epsilon} (y-x)^2 p(s, x; t, y) dy = b^2(s, x).$$

Alternatively:

$$a(s, x) = \lim_{t \rightarrow s^+} \frac{1}{t-s} E(X(t) - X(s) | X(s) = x),$$

$$b^2(s, x) = \lim_{t \rightarrow s^+} \frac{1}{t-s} E((X(t) - X(s))^2 | X(s) = x).$$

a : drift coefficient, b : diffusion coefficient.

Wiener: $(a, b) = (0, 1)$, O-U: $(a, b) = (-\gamma x, \sqrt{2\gamma})$. (Calculation in next part)

• Kolmogorov equation:

Forward:

$$p_t + (a(t, y)p)_y = \frac{1}{2}(b^2(t, y)p)_{yy},$$

Backward:

$$p_s + a(s, x)p_x = -\frac{1}{2}b^2(s, x)p_{xx}.$$

Forward equation is also called Fokker-Planck equation.

3 Calculating Drift and Diffusion

For Brownian motion, using independence of increment, we see that:

$$E(X(t) - X(s) | X(s) = x) = 0,$$

so

$$a(s, x) = \lim_{t \rightarrow s^+} \frac{1}{t-s} E(X(t) - X(s) | X(s) = x) = 0;$$

$$E((X(t) - X(s))^2 | X(s) = x) = t - s,$$

$$b^2(s, x) = \lim_{t \rightarrow s^+} \frac{1}{t-s} E((X(t) - X(s))^2 | X(s) = x) = 1.$$

Hence by Gaussian process:

$$p(s, x; t, y) = \frac{1}{\sqrt{2\pi(t-s)}} \exp\left\{-\frac{(y-x)^2}{2(t-s)}\right\}.$$

implying that BM satisfies:

$$E(|X(t) - X(s)|^4) = 3|t-s|^2, \quad \forall s, t \geq 0.$$

O-U:

Fact: $\tau \geq 0$, $X(t+\tau) - e^{-\gamma t}X(\tau)$ is independent of $(\omega : X(s), s \leq \tau)$.

Covariance ($s \leq \tau$):

$$\begin{aligned} E[(X(t+\tau) - e^{-\gamma t}X(\tau))X(s)] &= C(s, t+\tau) - e^{-\gamma t}C(s, \tau) \\ &= e^{-\gamma|t+\tau-s|} - e^{-\gamma t - \gamma|\tau-s|} \\ &= 0. \end{aligned} \tag{3.1}$$

At the same time, X is a Gaussian process.

To find transition probability:

$$\begin{aligned} P(X(t+\tau) \in A | X(\tau) = x) &= \\ P(X(t+\tau) - e^{-\gamma t}X(\tau) \in A - e^{-\gamma t}X(\tau) | X(\tau) = x) &= \\ = P(X(t+\tau) - e^{-\gamma t}X(\tau) \in A - e^{-\gamma t}x). \end{aligned} \tag{3.2}$$

R.V. $X(t+\tau) - e^{-\gamma t}X(\tau)$ is mean zero, Gaussian, and has variance:

$$\begin{aligned} E[(X(t+\tau) - e^{-\gamma t}X(\tau))^2] &= \\ E[(X(t+\tau) - e^{-\gamma t}X(\tau))X(t+\tau)] &= \\ = 1 - e^{-2\gamma t}. \end{aligned} \tag{3.3}$$

(3.2) and (3.3) imply:

$$p(s, x; t, y) = \frac{1}{\sqrt{2\pi(1 - e^{-2\gamma(t-s)})}} \exp\left\{-\frac{(y - xe^{-\gamma(t-s)})^2}{2(1 - e^{-2\gamma(t-s)})}\right\}.$$

Calculate drift:

$$\begin{aligned} E(X(t) - X(s) | X(s) = x) &= E[X(t) - e^{-\gamma|t-s|}X(s) \\ &+ e^{-\gamma|t-s|}X(s) - X(s) | X(s) = x] \\ &= (e^{-\gamma|t-s|} - 1)x, \end{aligned} \tag{3.4}$$

$$a(s, x) = -\gamma x.$$

Calculate diffusion:

$$\begin{aligned} E((X(t) - X(s))^2 | X(s) = x) &= E[(X(t) - e^{-\gamma|t-s|}X(s) \\ &+ (e^{-\gamma(t-s)} - 1)x)^2] \\ &= 1 - e^{-2\gamma(t-s)} + (e^{-\gamma(t-s)} - 1)x^2, \end{aligned} \quad (3.5)$$

$$b^2(s, x) = 2\gamma.$$

Over small time interval $[s, t]$, using drift-diffusion information, we see that O-U is related to BM as (to leading order):

$$X(t) - X(s) = -\gamma X(s)(t - s) + \sqrt{2\gamma}(W(t) - W(s)),$$

where $W(t)$ denotes BM; or in differential form:

$$dX = -\gamma X dt + \sqrt{2\gamma} dW,$$

The term $-\gamma X dt$ physically means damping.

4 From Random Walk to Brownian Motion

Divide time interval $[0, 1]$ into N equal length subintervals $[t_i, t_{i+1}]$, $i = 0, 1, \dots, N$. Consider a walker making steps $\pm\sqrt{\delta t}$, $\delta t = 1/N$ with probability $1/2$ each, starting from $x = 0$. In n steps, the walker's location is:

$$S_N(t_n) = \sqrt{\delta t} \sum_{i=1}^n X_i, \quad (4.6)$$

where X_i are independent two point r.v's taking ± 1 with equal probability. Define a piecewise continuous function:

$$S_N(t) = S_N(t_n), \quad t \in [t_n, t_{n+1}], \quad n \leq N - 1.$$

S_N has independent increment $X_1\sqrt{\delta t}$, $X_2\sqrt{\delta t}$ etc for given subintervals, and in the limit $N \rightarrow \infty$ tends to a process with independent increment. Moreover:

$$E(S_N) = 0, \quad Var(S_N(t)) = \delta t \left[\frac{t}{\delta t} \right].$$

In the limit $N \rightarrow \infty$, $Var(S_N(t)) \rightarrow t$. Applying Central Limit Theorem, $\forall t$, the approximate process $S_N(t)$ converges in law to a process with independent increment, zero mean, variance t , and Gaussian. So it is a BM.

Using $S_N(t)$ is a way to *numerically construct BM* as well. The X_i 's are generated from $U(0, 1)$ as: $X_i = 1$ if $U \in [0, 1/2]$; $X_i = -1$, if $U \in (1/2, 1]$. A shortcut is to replace two point X_i 's by i.i.d unit Gaussian r.v's. Try the 2 line Matlab code to generate a BM sample path:

```
randn('state',0); N=1e4; dt=1/N;
```

```
w=sqrt(dt)*cumsum([0;randn(N,1)]); plot([0:dt:1],w);
```

`cumsum` is a fast summation on vector input. Change the state number from 0 to 10 (or a larger number if you are having fun !) to see different sample paths (see Figure 1).




Figure 1: Four Sample Paths of Numerical Approximation of Brownian Motion on $[0,1]$.

BM sample path is almost surely continuous. Kolmogorov criterion:

$$E(|X(t) - X(s)|^a) \leq C|t - s|^{1+b},$$

for a, b, C positive. For BM, $a = 4, b = 1, C = 3$.

Kolmogorov Continuity Theorem:

Let (S, d) be some complete metric space, and let $X : [0, +\infty) \times \Omega \rightarrow S$ be a stochastic process. Suppose that for all times $T > 0$, there exist positive constants α, β, K such that

$\mathbb{E}[d(X_t, X_s)^\alpha] \leq K|t-s|^{1+\beta}$ for all $0 \leq s, t \leq T$. Then there exists a modification \tilde{X} of X that is a continuous process,

i.e. a process $\tilde{X} : [0, +\infty) \times \Omega \rightarrow S$ such that

- \tilde{X} is sample-continuous;
- for every time $t \geq 0$, $\mathbb{P}(X_t = \tilde{X}_t) = 1$ (modification)

Furthermore, the paths of \tilde{X} are locally γ -Hölder-continuous for every $0 < \gamma < \frac{\beta}{\alpha}$.

5 BM via Random Fourier Series

Let $Y_i, i = 1, 2, \dots$ be independent, $N(0, 1)$; $\varphi_k(t)$ be defined on closed interval I such that

$$\sum_{k=1}^{\infty} |\varphi_k(t)|^2 < \infty, \quad t \in I.$$

$$\sum_{k=1}^{\infty} E|\varphi_k(t)Y_k|^2 = \sum_{k=1}^{\infty} |\varphi_k(t)|^2 < \infty,$$

so:

$$Z(t) = \sum_{k=1}^{\infty} \varphi_k(t)Y_k,$$

converges a.s. in t .

$E[Z(t)] = 0$, covariance:

$$C(t, s) = \sum_{k=1}^{\infty} \varphi_k(t) \varphi_k(s),$$

To match Z with BM, require:

$$\min(t, s) = \sum_{k=1}^{\infty} \varphi_k(t) \varphi_k(s).$$

Let $I = [0, \pi]$, fact:

$$\min(t, s) = \frac{ts}{\pi} + \frac{2}{\pi} \sum_{k \geq 1} \frac{\sin kt \sin ks}{k^2}.$$

$$W(t) = BM = \frac{t}{\sqrt{\pi}}Y_0 + \sqrt{\frac{2}{\pi}} \sum_{k \geq 1} \frac{\sin kt}{k} Y_k, \quad (5.7)$$

$t \in [0, \pi]$, $W(t)$ standard BM. By truncating the random Fourier series, we have a second way to generate BM.

6 Spectral Representation

Consider stationary process, e.g. O-U. Covariance $C(t, s) = C(t - s)$, $C(\cdot)$ even function, and: $\forall \{a_i\} \subset R$,

$$\sum_{k,j} a_k a_j C(t_k - t_j) = E(|\sum_k a_k X(t_k)|^2) \geq 0,$$

$C(\cdot)$ is nonnegative definite and symmetric. Bochner theorem:

$$C(t) = \int_{-\infty}^{\infty} e^{2\pi i t \lambda} dF(\lambda), \quad (6.8)$$

$F(\lambda)$ is nondecreasing, right continuous, $F(+\infty) - F(-\infty) = C(0)$. We call F *spectral distribution function* of process $X(t)$, $F'(\lambda)$ *spectral density*.

$$F'(\lambda) = \int_{R^1} C(s) e^{-2\pi i \lambda s} ds = \int_{R^1} C(s) \cos(2\pi \lambda s) ds. \quad (6.9)$$

Just like finding a random Fourier series for BM from its covariance, one can construct a random Fourier integral for $X(t)$:

Let $Z(\lambda)$ be a process with orthogonal increments:

$$E[(Z(a) - Z(b))(Z(a') - Z(b'))] = 0,$$

if $(a, b) \cap (a', b')$ empty, and

$$E[(Z(a) - Z(b))^2] = F(a) - F(b),$$

Now:

$$X(t) = \int_{-\infty}^{\infty} e^{2\pi i t \lambda} dZ(\lambda). \quad (6.10)$$

The random integral $\int g(\lambda) dZ(\lambda)$ is defined as mean square limit of finite Stieltjes sum:

$$\sum g_k [Z(\lambda_k) - Z(\lambda_{k-1})],$$

if $g \in L^2(dF)$.

Examples of Spectral Densities:

(1) O-U: $C(s) = e^{-\gamma|s|}$, taking Fourier transform (6.9):

$$F'(\lambda) = \frac{2\gamma}{\gamma^2 + 4\pi^2 \lambda^2}.$$

(2) Gaussian white noise: $F'(\lambda) = \text{constant}$, $C(s) = c_0 \delta(s)$.

Approximated by:

$$X_h(t) = (W(t+h) - W(t))/h,$$

small $h > 0$. Process X^h has covariance and spectral density:

$$C_h(s, t) = \frac{1}{h} \max(0, 1 - |t - s|/h),$$

$$F'_h(\lambda) = \sin^2(2\pi\lambda h)/(\pi\lambda h)^2,$$

broad band spectrum, X_h called colored noise. In the limit $h \rightarrow 0$, C_h converges to delta function, X_h converges in some weak sense to white noise.